

This article was downloaded by: [University of Limerick]

On: 21 May 2013, At: 14:48

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Ergonomics

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/terg20>

Lumbar posture and trunk muscle activation during a typing task when sitting on a novel dynamic ergonomic chair

Kieran O'Sullivan^a, Raymond McCarthy^a, Alison White^a, Leonard O'Sullivan^{a,b} & Wim Dankaerts^c

^a Department of Clinical Therapies, University of Limerick, Limerick, Ireland

^b Department of Design and Manufacturing Technology, University of Limerick, Limerick, Ireland

^c Department of Rehabilitation Sciences, Catholic University, Leuven, Belgium

Published online: 25 Sep 2012.

To cite this article: Kieran O'Sullivan, Raymond McCarthy, Alison White, Leonard O'Sullivan & Wim Dankaerts (2012): Lumbar posture and trunk muscle activation during a typing task when sitting on a novel dynamic ergonomic chair, *Ergonomics*, 55:12, 1586-1595

To link to this article: <http://dx.doi.org/10.1080/00140139.2012.721521>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Lumbar posture and trunk muscle activation during a typing task when sitting on a novel dynamic ergonomic chair

Kieran O'Sullivan^{a*}, Raymond McCarthy^a, Alison White^a, Leonard O'Sullivan^{a,b} and Wim Dankaerts^c

^aDepartment of Clinical Therapies, University of Limerick, Limerick, Ireland; ^bDepartment of Design and Manufacturing Technology, University of Limerick, Limerick, Ireland; ^cDepartment of Rehabilitation Sciences, Catholic University, Leuven, Belgium

(Received 17 September 2011; final version received 10 August 2012)

Low back pain (LBP) is a common musculoskeletal disorder and prolonged sitting often aggravates LBP. A novel dynamic ergonomic chair ('Back App'), which facilitates less hip flexion while sitting on an unstable base has been developed. This study compared lumbar posture and trunk muscle activation on this novel chair with a standard backless office chair. Twelve painfree participants completed a typing task on both chairs. Lumbar posture and trunk muscle activation were collected simultaneously and were analysed using paired *t*-tests. Sitting on the novel dynamic chair significantly ($p < 0.05$) reduced both lumbar flexion and the activation of one back muscle (Iliocostalis Lumborum pars Thoracis). The discomfort experienced was mild and was similar ($p > 0.05$) between chairs. Maintaining lordosis with less muscle activation during prolonged sitting could reduce the fatigue associated with upright sitting postures. Studies with longer sitting durations, and in people with LBP, are required.

Practitioner Summary: Sitting on a novel dynamic chair resulted in less lumbar flexion and less back muscle activation than sitting on a standard backless office chair during a typing task among pain-free participants. Facilitating lordotic sitting with less muscle activation may reduce the fatigue and discomfort often associated with lordotic sitting postures.

Keywords: posture; back pain; ergonomics; lordosis; sitting

1. Introduction

Low back pain (LBP) is a very common and costly disorder (Dagenais *et al.* 2008). While it is commonly thought that prolonged sitting may be a risk factor for the development of LBP, recent systematic reviews demonstrate that sitting duration alone is not linked to the onset of LBP (Lis *et al.* 2007, Bakker *et al.* 2009, Roffey *et al.* 2010). As a result, other sitting parameters are of interest, including the chair used, the amount of spinal motion in sitting and the spinal curvature during sitting.

Although sitting is associated with greater lumbar flexion than standing (Scannell and McGill 2003, De Carvalho *et al.* 2010), there is no consensus regarding what is an optimal sitting posture (Claus *et al.* 2009b, O'Sullivan *et al.* 2010). Both upright (Vergara and Page 2002) and slump (Womersley and May 2006) sitting postures can be provocative for patients with LBP (O'Sullivan 2000). Slump sitting results in a 'flexion relaxation' response of the paraspinal muscles, potentially increasing the strain on passive spinal structures (Andersson *et al.* 1996, O'Sullivan *et al.* 2006b). In contrast, more lordotic sitting postures are associated with greater trunk muscle activity with the potential cost of increased trunk muscle fatigue (O'Sullivan *et al.* 2006a, Claus *et al.* 2009a,b).

Prolonged low level muscle activity has been linked to pain in other muscle groups (Westgaard and DeLuca 1999). More neutral sitting postures, involving slight lumbar lordosis and a relaxed thorax, have been recommended to facilitate activation of key trunk muscles without excessive compressive spinal load (O'Sullivan *et al.* 2010). Since posture is influenced by a wide range of factors including genetics (Seah *et al.* 2011), gender (Dunk and Callaghan 2005, Smith *et al.* 2010), body mass index (Smith *et al.* 2011) and psychological factors (O'Sullivan *et al.* 2011a), the best sitting posture may need to consider these factors as well as individual variations in specific aggravating/easing factors (Dankaerts *et al.* 2009).

Adjustable chairs that reduce hip flexion to promote lordotic lumbar postures have been associated with decreased lumbar muscle tension and improved muscle strength over a two-year period, but did not decrease LBP incidence (Koskelo *et al.* 2007). Using a different saddle chair promoted less hip flexion in pain-free subjects performing typing tasks whilst sitting for 2 h (Gadge and Innes 2007). Interestingly, while the saddle seat was

*Corresponding author. Email: kieran.osullivan@ul.ie

associated with significantly less lumbar discomfort, it was associated with significantly more hip/buttock discomfort (Gadge and Innes 2007). Another study (Gale *et al.* 1989) showed that the same saddle chair increased lumbar lordosis during a seated work task, but this study did not investigate the effect of this altered posture on muscle activation or levels of discomfort. Therefore, while there is some evidence that such saddle chairs may reduce hip and lumbar flexion, the effect of such seats on trunk muscle activity and sitting discomfort is unclear. This is important considering there is evidence that other ergonomic approaches such as kneeling-chairs which effectively enhance lordosis (Bennett *et al.* 1989, Bettany-Saltikov *et al.* 2008), do not alter muscle activation (Bennett *et al.* 1989), or might actually be associated with both increased back muscle activation and increased discomfort (Lander *et al.* 1987).

Another relevant factor may be the degree of spinal movement during sitting. During prolonged sitting, people tend to choose a varied rather than a single comfortable position (Callaghan and McGill 2001), thereby frequently changing the postural load. It has been hypothesised that dynamic sitting on an unstable base of support may facilitate spinal motion and help prevent muscle fatigue via alternating motor unit activation (van Dieen *et al.* 2001). Dynamic sitting has been proposed to reduce spinal shrinkage (van Deursen *et al.* 1999, van Dieen *et al.* 2001). However, the evidence from most studies suggest dynamic sitting results in little or no change in lumbar posture, trunk muscle activation or discomfort (van Dieen *et al.* 2001, Gregory *et al.* 2006, McGill *et al.* 2006, O'Sullivan *et al.* 2006c, Kingma and van Dieen 2009). To date, a comparison of a standardised seated office task with simultaneous monitoring of lumbar posture, trunk muscle activation and levels of discomfort has not been conducted.

The 'Back App' is a commercially available ergonomic chair that incorporates both a saddle-design to reduce hip flexion, as well as an unstable base of support. Both the chair height and the degree of instability can be adjusted. It has the potential to facilitate less flexed lumbar postures and greater spinal micro-movement, although there are currently no published data available on its effect on lumbar posture and trunk muscle activity. Therefore, the aims of this study were to compare the lumbar posture and trunk muscle activation of this novel dynamic ergonomic chair with a standard backless office chair during a typing task. The study hypothesis was that the 'Back App' would be associated with less lumbar flexion in sitting, without an increase in trunk muscle activation.

2. Methods

2.1. Study design

A single session, repeated measures, crossover study was conducted. The dependent variables were lumbar posture, trunk muscle activation and discomfort. The independent variable was chair type ('Back App' and standard chair). All participants completed the same protocol apart from the order in which they sat on the chairs, which they randomly selected from a sealed opaque envelope. Ethical approval was obtained from the local university Research Ethics Committee.

2.2. Participants

Twelve (7 F, 5 M) pain-free participants were recruited from the local community. All participants provided written informed consent. Participants were aged >18 years, were not pregnant, had no LBP in the last two years, no previous spinal surgery, no current pain medications, had not undertaken previous postural control training, and could speak/understand English. Participants' mean(SD) age was 23.3(3.6) years, height was 169.5(5.7) cm, mass was 65.9(10.2) kg and body mass index was 22.9(3.2) kg/m².

2.3. Instrumentation

2.3.1. Kinematics

Postural data were collected using a wireless posture monitor ('BodyGuard') (Figure 1). The 'BodyGuard' (Sels Instruments, Belgium) incorporates a strain gauge that provides information about the relative distance between anatomical landmarks, estimating flexion/extension range of the lumbar spine by the degree of strain gauge elongation. Elongation of the strain gauge alters its internal resistance and therefore the voltage of the signal. This alteration in voltage occurs in a linear manner in response to elongation. Therefore, the voltage output is directly related to the length (flexion vs. extension) of the strain gauge. Based on the elongation of the strain gauge, lower lumbar spine sagittal plane posture is expressed as a percentage of range of motion (ROM). Therefore, the degree of




Figure 1. The BodyGuard posture monitor.

spinal flexion/extension is expressed relative to a referenced ROM, for example, total lumbar flexion ROM, rather than being expressed in degrees (O'Sullivan *et al.* 2010). This reflects the clinical assessment of patients, where sitting posture is often considered relative to individual ROM. Calculation of posture relative to ROM has been used in previous spinal posture research (Edmondston *et al.* 2007). It is also similar to electromyography normalisation of muscle activity relative to maximal or sub-maximal voluntary contraction (Dankaerts *et al.* 2006b). This posture monitor has been shown to have very good reliability (ICC > 0.84) (O'Sullivan *et al.* 2011b) and validity (Spearman's correlation > 0.88) (O'Sullivan *et al.* 2012) for the measurement of lumbar posture.

Recent research suggests that the upper and lower lumbar spine regions demonstrate functional independence, with the lower lumbar spine being the most common area for subjects to report non-specific chronic low back pain (NSCLBP) (Dankaerts *et al.* 2006a), and the area demonstrating the greatest postural differences among LBP subjects (Dankaerts *et al.* 2006b, Mitchell *et al.* 2008). Consequently, a strain gauge was positioned directly over the spine at the spinal levels of L3 and S2, after manual palpation of these spinal levels in a slightly flexed sitting posture. Participants then performed maximal lumbar ROM to ensure the device was securely attached. To calibrate the posture monitor, manual and verbal facilitation were used to guide subjects into a fully lordotic sitting posture which was set as 0% of their lumbar ROM, and then into a fully flexed sitting posture which was set as 100% of their lumbar ROM (O'Sullivan *et al.* 2010). This was repeated five times, to obtain a representative ROM value. Postural data were recorded continuously in real-time at 1 Hz.

2.3.2. Trunk muscle activation

The activation of six trunk muscles was analysed using surface electromyography (sEMG). A Motion Lab Systems MA-300 multi-channel EMG system (Motion Lab Systems Inc., Baton Rouge, Louisiana, USA) collected sEMG data using bipolar, pre-amplified, circular electrodes 144 mm² in size, with a fixed inter-electrode distance of 18 mm. The sampling rate was 1000 Hz per channel, with a bandwidth of 0–500Hz and a gain of 2000. The common mode rejection ratio was > 100 dB at 60 Hz. Three abdominal and three back muscles of the right hand side of the trunk were analysed, after preliminary testing had demonstrated no significant difference between right and left sides in pain-free controls during this task. The skin was prepared for electrode placement by abrading the skin with fine sandpaper, shaving any hair and cleansing the skin with isopropyl alcohol solution to reduce skin impedance, in line with agreed international recommendations (Hermens *et al.* 2000). Pairs of surface electrodes were positioned parallel to the muscle fibre direction of each individual muscle and secured with clear adhesive tape. The muscles studied were superficial lumbar multifidus (LM) (L5 level, parallel to a line connecting the posterior superior iliac spine and L1–L2 interspinous space); iliocostalis lumborum pars thoracis (ICLT) (level of L1 spinous process, midway between the midline and lateral aspect of the participant's body); thoracic erector spinae (TES) (5 cm lateral to the T9 spinous process); external oblique (EO) (just below the rib cage, along a line connecting the most inferior costal margin and the contralateral pubic tubercle); internal oblique (IO) (1 cm medial to the anterior superior iliac spine) and rectus abdominis (RA) (1 cm above the umbilicus and 2 cm lateral to midline). These

electrode placements were consistent with previous research (O'Sullivan *et al.* 2006a). A common earth electrode was placed over the ulnar styloid. Correct location of the electrodes was visually confirmed by examining the sEMG output while applying manual resistance. Electromyography (EMG) data were normalised to maximum voluntary isometric contraction (MVIC). To generate MVIC for the abdominal muscles, three variations of a sit-up were used, similar to previous research (O'Sullivan *et al.* 2006a). One normalisation technique was used for all three back muscles, similar to previous research (O'Sullivan *et al.* 2006a). The middle 3 s of amplitude normalised EMG data, from the 5 s testing period, were analysed. The highest generated contraction from any of the three abdominal tests was taken as the MVIC for each specific abdominal muscle, and the highest generated MVIC from three repetitions of the back muscle test was taken for each specific back muscle (O'Sullivan *et al.* 2006a). To avoid fatigue contraction time for all MVIC trials was 5 s duration (Soderberg and Knutson 2000) and a 3 min rest was given between trials (McLean *et al.* 2003).

2.3.3. Chairs

The 'Back App' chair (Figure 2) facilitates dynamic sitting through an unstable ball positioned at its base, whose prominence can be altered to vary the degree of motion allowed, and thereby the postural challenge. For testing, the degree of motion allowed on the 'Back App' chair was standardised at the 'green zone', which involves a mild degree of movement. The standard office chair (Figure 3) was adjustable, backless and had wheels. Participants were instructed to 'sit as you normally would' while on the chair, and simply to maintain their balance while sitting on the 'Back App'.




Figure 2. Sitting on the 'Back App' chair.




Figure 3. Sitting on the standard office chair.

2.3.4. Discomfort

Participant discomfort was rated numerically during the typing task using the body part discomfort scale (BPDS) (Fenety and Walker 2002). This involved participants rating discomfort across each of 12 body areas from 0 (no discomfort) to 5 (intolerable discomfort) (Fenety and Walker 2002). Low back discomfort, as well as overall body discomfort using the mean discomfort of the 12 body parts, was recorded (Fenety and Walker 2002).

2.4. Procedure

2.4.1. Workstation set-up

A simulated workstation was created. An adjustable height desk was elevated until it reached the underneath of the elbow, to allow a 90° elbow angle in line with the trunk (Kingma and van Dieen 2009). Following this, the desk was positioned in line with the radial styloid process. Participants distance from the desk was standardised as their greater trochanter being 30 cm from the desk. Goniometry was used to measure both hip and knee angles. The standard chair was adjusted to allow an angle of 90° for both the hips and knees with the feet placed firmly on the floor (Figure 1), while the 'Back App' was adjusted to allow a 125° hip angle with the feet placed on the footplate (Figure 2). Participants were blinded as to when all posture and sEMG measurements were recorded. After chair order was assigned, participants completed the typing task, with a 1-min break given while changing between chairs.

2.4.2. Typing task

A laptop was placed 10 cm from the edge of the desk. Participants typed the same piece of literature, placed on a stand to the side of the laptop, for 10 min on each chair. Electromyography (EMG) data were recorded on three occasions (after three, six and nine min) for 5 s duration, similar to previous research (McGill *et al.* 2006). At time intervals 0, 5 and 10 min of the typing task, participants rated discomfort using the BPDS. To control for variations

in baseline discomfort level and order of testing, the progression of discomfort over time (discomfort after 10 min – discomfort at baseline) was used to determine the discomfort associated with each sitting condition.

2.5. Data analysis

Posture did not change significantly over time during the typing task on both chairs. Consequently, a single value for average (mean) posture and variation in posture (SD) was calculated for each participant for each chair.

All raw EMG files were visually inspected. The middle 3 s of raw EMG data from all testing periods was processed using a root-mean-square (RMS) algorithm, and expressed as %MVIC. For the typing task, the mean of the three values was used, as there was minimal variance between them, in line with previous studies who found no significant effect of time on trunk muscle activity in short duration sitting tasks (McGill *et al.* 2006).

Data were analysed using SPSS 18.0. For all statistical tests, significance was set at $p < 0.05$. Data were tested for normality using the Shapiro–Wilks test. Paired *t*-tests were used to compare both posture (mean and variation) and trunk muscle activity (mean and variation) during the typing task. A Wilcoxon signed-ranks test was used to compare mean discomfort on both chairs.

3. Results

The overall results for lumbar posture and trunk muscle activation on both chairs during the typing task are illustrated in Table 1.

3.1. Posture

There was significantly less ($p = 0.002$) lumbar flexion when sitting on the ‘Back App’; however, the variation in posture was not significantly different ($p > 0.05$) (Table 1).

3.2. Trunk muscle activation

Despite ‘Back App’ sitting involving less flexion, mean ICLT activity was significantly lower ($p = 0.015$) on the ‘Back App’ chair than on the standard chair. None of the other five trunk muscles measured demonstrated a significant difference in activity (Figure 4). There was no difference in the variation of trunk muscle activity between the standard backless office chair and the ‘Back App’ chair (all $p > 0.05$).

3.3. Discomfort

The level of both low back discomfort and overall body discomfort experienced during the typing task was mild, and did not differ significantly ($p > 0.05$) between the standard backless office chair and the ‘Back App’ chair.

Table 1. Mean(SD) trunk muscle activation and lumbar posture during a typing task on both a standard office chair and the ‘Back App’ chair.

	Standard chair	Back App	<i>p</i> -value
EO	5.6(3.8)	4.0(1.4)	0.342
IO	7.6(6.0)	7.3(7.5)	0.951
RA	6.4(6.8)	3.9(2.6)	0.107
TES	12.0(5.8)	10.2(5.5)	0.278
ICLT*	11.9(4.8)	8.3(3.9)	0.015
LM	7.3(5.5)	5.3(3.2)	0.160
% Flexion (Mean) [†]	70.13	46.34	0.002
% Flexion (Variation)	6.72	4.98	0.328

Note: EO – external oblique; IO – transverse fibres of internal oblique; RA – rectus abdominis; TES – thoracic erector spinae; ICLT – iliocostalis lumborum pars thoracis; LM – superficial fibres of lumbar multifidus; All muscle activation expressed as %MVIC (Maximum Voluntary Isometric Contraction); All posture expressed as %flexion range of motion (ROM); *Significantly less activation on ‘Back App’; [†]Significantly less flexion on ‘Back App’.




Figure 4. Mean(SD) trunk muscle activation during a typing task on both a standard office chair and the 'Back App' chair. Note: %MVIC – percentage of Maximum Voluntary Isometric Contraction; EO – external oblique; IO – transverse fibres of internal oblique; RA – rectus abdominis; TES – thoracic erector spinae; ICLT – iliocostalis lumborum pars thoracis; LM – superficial fibres of lumbar multifidus; *Statistically significant difference.

4. Discussion

The results indicate that during a brief typing task, pain-free participants sat in less lumbar flexion, and displayed less lumbar muscle activation, when sitting on a novel dynamic ergonomic chair compared to a standard backless office chair. This change in lumbar posture and lower paraspinal muscle activation was achieved without altering mean overall body discomfort.

No previous study had examined the effect of a dynamic saddle chair similar to the 'Back App' on lumbar posture or trunk muscle activity. The current results are consistent with data demonstrating that static saddle seats (Gale *et al.* 1989, Koskelo *et al.* 2007) are associated with increased lumbar lordosis. However, the existing data on the effects of saddle chairs on trunk muscle activation are less clear. Consistent with our results, one study (Koskelo *et al.* 2007) demonstrated reduced muscle activation at two year follow-up among students who used adjustable saddle chairs in the intervening two years. However, another study (Bennett *et al.* 1989) observed no difference in muscle activation when sitting on these adjustable chairs, possibly due to using chairs with a backrest for comparison, since lumbar supports and increased seat back inclination angle also reduce paraspinal muscle activation (Magnusson 1998). In contrast to the current results, it appears that other ergonomic methods of increasing lumbar lordosis such as kneeler chairs increase lumbar lordosis but at the cost of increasing paraspinal muscle activation and discomfort (Lander *et al.* 1987, Bennett *et al.* 1989).

The existing research on the influence of dynamic sitting is contradictory. It appears to increase the overall amount of spinal motion in sitting (O'Sullivan *et al.* 2006c, Kingma and van Dieen 2009), but the effects on height of the spinal column (spinal shrinkage) are unclear (van Dieen *et al.* 2001, Kingma and van Dieen 2009). Most of the evidence suggests that dynamic sitting is associated with no change in mean spinal posture (van Dieen *et al.* 2001, McGill *et al.* 2006, O'Sullivan *et al.* 2006c) although it may increase anterior pelvic tilt (Gregory *et al.* 2006). Similarly, most research suggests dynamic sitting does not change (van Dieen *et al.* 2001, McGill *et al.* 2006, O'Sullivan *et al.* 2006c) or in fact increases (Gregory *et al.* 2006, Kingma and van Dieen 2009) trunk muscle activation, in contrast to the trend towards reduced lower paraspinal muscle activation seen in the current study. Finally, dynamic sitting may be associated with increased discomfort (Gregory *et al.* 2006), unlike the current study where the 'Back App' was associated with similar levels of discomfort. Overall, since most previous research on dynamic sitting suggests no major differences in posture or muscle activation (van Dieen *et al.* 2001, Gregory *et al.* 2006, McGill *et al.* 2006, O'Sullivan *et al.* 2006c, Kingma and van Dieen 2009), it is likely that the changes seen in posture and muscle activity while sitting on the 'Back App' are primarily related to the reduction in hip flexion. Specifically, the 35° difference in hip angle between the chairs may be critical. As a result, if dynamic sitting has a benefit, it may be in preventing static loading of spinal tissues which may be relevant in LBP (Vergara and Page 2002) rather than as a means of altering trunk posture or trunk muscle activation (van Deursen *et al.* 1999, van Dieen *et al.* 2001; O'Sullivan *et al.* 2006c). The lack of any significant difference in abdominal muscle activation on the chairs suggests that the novel dynamic ergonomic chair has minimal effect on abdominal muscle activation.

Sitting is associated with increased spinal flexion compared to standing (Dunk *et al.* 2009, De Carvalho *et al.* 2010). It has been observed that while sitting on traditional chairs, paraspinal muscle activation is closely related to seated lumbar curvature (O'Sullivan *et al.* 2006a). For example, actively increasing anterior pelvic tilt and lumbar

lordosis are associated with increased paraspinal muscle activation (O'Sullivan *et al.* 2006a, Claus *et al.* 2009a). However, in this study, back muscle activation, especially ICLT, was reduced even with an increase of lumbar lordosis while sitting on the 'Back App'. This suggests that the 'Back App' chair passively assists lumbar lordosis, without requiring an increased level of paraspinal muscle activation. Considering suggestions that sustaining lordotic sitting postures may place an unsustainable stress on the back muscles and contribute to pain and fatigue (Claus *et al.* 2009b), this is potentially advantageous in moderating spinal loads. The values for trunk muscle activation in the current study appear representative of previous research (O'Sullivan *et al.* 2006a, Claus *et al.* 2009a) and are minimal, with mean trunk muscle activation ranges from 4 to 12% MVIC.

Considering that the 'Back App' is likely to alter lumbar lordosis by reducing posterior pelvic rotation, it is not surprising that the lower paraspinal muscles (ICLT and LM) are most influenced by its use. While sitting posture also influences the activation of the abdominal muscles and TES, these muscles are not as closely affected by changes in pelvic rotation (O'Sullivan *et al.* 2006a, Claus *et al.* 2009a). While there is considerable evidence that the activation of muscles such as LM and transversus abdominis may be delayed in people with LBP (Hodges 2001, MacDonald *et al.* 2009), there is also evidence that overall amplitude of trunk muscle activation may be increased in some subjects with LBP (Dankaerts *et al.* 2006a).

Sustained sitting can induce significant discomfort (Vergara and Page 2002), which can be alleviated by the use of backrests (Gale *et al.* 1989, Leivseth and Drerup 1997). Both chairs in the current study induced very low discomfort levels, which did not differ significantly. The short duration of testing, the use of a rest-break between the chairs, and the painfree nature of the participants may explain the low levels of discomfort reported. A previous study (Gadge and Innes 2007) reported that overall body and lumbar discomfort were slightly reduced during a typing task on a static saddle chair compared to a standard office chair, but that lower limb discomfort was increased on the saddle chair. A similar study (Koskelo *et al.* 2007) reported that students using adjustable height chairs were more comfortable than those using traditional chairs. In contrast, in another study (Gregory *et al.* 2006) overall body discomfort was significantly higher after a 1-h period of dynamic sitting compared to a standard office chair. However, this increased discomfort may be explained by the fact that in their study the exercise ball appears to have facilitated anterior pelvic tilt through increased paraspinal muscle activation, which differs significantly from the reduced paraspinal muscle activation observed in the current study when using the 'Back App'.

It has to be acknowledged that there are some limitations to the current study. This study involved only a small sample of pain-free participants, without an a priori power calculation of the sample size necessary to detect differences between the chairs. In addition, the duration of exposure was relatively short. Both the short duration and small sample size reduce the likelihood of finding significant differences between the two sitting conditions. Nevertheless, significant differences in posture and muscle activity were still evident. Differences in posture, muscle activation and particularly discomfort may be even more pronounced during longer sitting exposures in a larger sample of participants with LBP. Analysis of more dynamic seated tasks is warranted. Most standard office chairs have backrests which may also decrease the muscular effort and discomfort of sitting (Vergara and Page 2000), and comparison of the 'Back App' to a standard office chair with a backrest should be completed. However, we chose to compare to a chair without a backrest to initially examine the influence of an altered hip angle and a degree of instability without the possible influence of a backrest. Furthermore, backrest use is reduced in many office tasks such as typing (Vergara and Page 2000), diminishing the importance of chair backrests during the task examined in this study. The 'Back App' has the potential to vary the level of instability, and the effect of greater levels of instability is unclear. Future studies may consider the effect of 'Back App' sitting in occupational settings for a longer duration. Notwithstanding these limitations, the study had many strengths such as the use of a closely standardised functional task that replicates an office situation.

5. Conclusion

The use of a novel dynamic ergonomic chair facilitates a less flexed lumbar spine posture, while requiring less intense activation of the lower paraspinal muscles, especially ICLT, during a brief seated typing task. The degree of discomfort was low and similar on both chairs. It is likely that this effect is achieved by the reduced hip flexion passively facilitating anterior pelvic tilt and lumbar lordosis. The relative contribution of the dynamic sitting element is unclear. Maintaining lumbar lordosis with less intense muscle activation is potentially advantageous during prolonged sitting, as it could reduce the potential for fatigue and discomfort often associated with lordotic sitting postures. Future studies in subjects with LBP are warranted.

Acknowledgements

The study was supported by 'Back App', who provided the novel dynamic ergonomic chair, and part-funded the study. The lead author (KOS) is funded by a research fellowship from the Health Research Board of Ireland.

References

- Andersson, E.A., Oddsson, L.I., Grundstrom, H., Nilsson, J., and Thorstensson, A., 1996. EMG activities of the quadratus lumborum and erector spinae muscles during flexion-relaxation and other motor tasks. *Clinical Biomechanics*, 11, 392–400.
- Bakker, E.W., Verhagen, A.P., van Trijffel, E., Lucas, C., and Koes, B.W., 2009. Spinal mechanical load as a risk factor for low back pain: a systematic review of prospective cohort studies. *Spine*, 34, 281–293.
- Bennett, D.L., Gillis, D.K., Portney, L.G., Romanow, M., and Sanchez, A.S., 1989. Comparison of integrated electromyographic activity and lumbar curvature during standing and during sitting in three chairs. *Physical Therapy*, 69, 902.
- Bettany-Saltikov, J., Warren, J., and Jobson, M., 2008. Ergonomically designed kneeling chairs are they worth it?: comparison of sagittal lumbar curvature in two different seating postures. *Studies in Health Technology and Informatics*, 140, 103–106.
- Callaghan, J.P. and McGill, S.M., 2001. Low back joint loading and kinematics during standing and unsupported sitting. *Ergonomics*, 44, 280–294.
- Claus, A.P., Hides, J.A., Moseley, G.L., and Hodges, P.W., 2009a. Different ways to balance the spine: subtle changes in sagittal spinal curves affect regional muscle activity. *Spine*, 34, E208–E214.
- Claus, A., Hides, J., Moseley, G.L., and Hodges, P., 2009b. Is "ideal" sitting real?: measurement of spinal curves in four sitting postures. *Manual Therapy*, 14, 404–408.
- Dagenais, S., Caro, J., and Haldeman, S., 2008. A systematic review of low back pain cost of illness studies in the United States and internationally. *The Spine Journal*, 8, 8–20.
- Dankaerts, W., O'Sullivan, P., Burnett, A., and Straker, L., 2006a. Altered patterns of superficial trunk muscle activation during sitting in nonspecific chronic low back pain patients: importance of subclassification. *Spine*, 31, 2017–2023.
- Dankaerts, W., O'Sullivan, P., Burnett, A., and Straker, L., 2006b. Differences in sitting postures are associated with non-specific chronic low back pain disorders when sub-classified. *Spine*, 31, 698–704.
- Dankaerts, W., O'Sullivan, P., Burnett, A., Straker, L., Davey, P., and Gupta, R., 2009. Discriminating healthy controls and two clinical subgroups of nonspecific chronic low back pain patients using trunk muscle activation and lumbosacral kinematics of postures and movements: a statistical classification model. *Spine*, 34, 1610–1618.
- De Carvalho, D., Soave, D., Ross, K., and Callaghan, J., 2010. Lumbar spine and pelvic posture between standing and sitting: a radiologic investigation including reliability and repeatability of the lumbar lordosis measure. *Journal of Manipulative and Physiological Therapeutics*, 33, 48–55.
- Dunk, N.M. and Callaghan, J.P., 2005. Gender-based differences in postural responses to seated exposures. *Clinical Biomechanics*, 20, 1101–1110.
- Dunk, N., Kedgley, A., Jenkyn, T., and Callaghan, J., 2009. Evidence of a pelvis-driven flexion pattern: are the joints of the lower lumbar spine fully flexed in seated postures? *Clinical Biomechanics*, 24, 164–168.
- Edmondston, S., Chan, H., Chi Wing Ngai, G., Warren, M., Williams, J., Glennon, S., and Netto, K., 2007. Postural neck pain: an investigation of habitual sitting posture, perception of 'good' posture and cervicothoracic kinaesthesia. *Manual Therapy*, 12, 363–371.
- Fenety, A. and Walker, J.M., 2002. Short-term effects of workstation exercises on musculoskeletal discomfort and postural changes in seated video display unit workers. *Physical Therapy*, 82, 578–589.
- Gadge, K. and Innes, E., 2007. An investigation into the immediate effects on comfort, productivity and posture of the Bambach™ saddle seat and a standard office chair. *Work*, 29, 189–203.
- Gale, M., Feather, S., Jensen, S., and Coster, G., 1989. Study of a workseat designed to preserve lumbar lordosis. *Australian Occupational Therapy Journal*, 36, 92–99.
- Gregory, D.E., Dunk, N.M., and Callaghan, J.P., 2006. Stability ball versus office chair: comparison of muscle activation and lumbar spine posture during prolonged sitting. *Human Factors*, 48, 142–153.
- Hermens, H., Freriks, B., Disselhorst-Klug, C., and Rau, G., 2000. Development of recommendations for SEMG sensors and sensor placement procedures. *Journal of Electromyography and Kinesiology*, 10, 361–374.
- Hodges, P., 2001. Changes in motor planning of feedforward postural responses of the trunk muscles in low back pain. *Experimental Brain Research*, 141, 261–266.
- Kingma, I. and van Dieen, J.H., 2009. Static and dynamic postural loadings during computer work in females: sitting on an office chair versus sitting on an exercise ball. *Applied Ergonomics*, 40, 199–205.
- Koskelo, R., Vuorikari, K., and Hänninen, O., 2007. Sitting and standing postures are corrected by adjustable furniture with lowered muscle tension in high-school students. *Ergonomics*, 50, 1643–1656.
- Lander, C., Korbon, G., DeGood, D., and Rowlingson, J., 1987. The Balans chair and its semi-kneeling position: an ergonomic comparison with the conventional sitting position. *Spine*, 12, 269–272.
- Leivseth, G. and Drerup, B., 1997. Spinal shrinkage during work in a sitting posture compared to work in a standing posture. *Clinical Biomechanics*, 12, 409–418.
- Lis, A., Black, K., Korn, H., and Nordin, M., 2007. Association between sitting and occupational LBP. *European Spine Journal*, 16, 283–298.
- MacDonald, D., Moseley, G., and Hodges, P., 2009. Why do some patients keep hurting their back? Evidence of ongoing back muscle dysfunction during remission from recurrent back pain. *Pain*, 142, 183–188.

- Magnusson, M., 1998. A review of the biomechanics and epidemiology of working postures. *Journal of Sound and Vibration*, 215, 965–976.
- McGill, S.M., Kavcic, N.S., and Harvey, E., 2006. Sitting on a chair or an exercise ball: various perspectives to guide decision making. *Clinical Biomechanics*, 21, 353–360.
- McLean, L., Chislett, M., Keith, M., Murphy, M., and Walton, P., 2003. The effect of head position, electrode site, movement and smoothing window in the determination of a reliable maximum voluntary activation of the upper trapezius muscle. *Journal of Electromyography and Kinesiology*, 13, 169–180.
- Mitchell, T., O'Sullivan, P., Burnett, A.F., Straker, L., and Smith, A., 2008. Regional differences in lumbar spinal posture and the influence of low back pain. *BMC Musculoskeletal Disorders*, 9, 152.
- O'Sullivan, P.B., 2000. Masterclass. Lumbar segmental 'instability': clinical presentation and specific stabilizing exercise management. *Manual Therapy*, 5, 2–12.
- O'Sullivan, P., Dankaerts, W., Burnett, A., Farrell, G., Jefford, E., Naylor, C., and O'Sullivan, K., 2006a. Effect of different upright sitting postures on spinal-pelvic curvature and trunk muscle activation in a pain-free population. *Spine*, 31, E707–E712.
- O'Sullivan, P., Dankaerts, W., Burnett, A., Chen, D., Booth, R., Carlsen, C., and Schultz, A., 2006b. Evaluation of the flexion relaxation phenomenon of the trunk muscles in sitting. *Spine*, 31, 2009–2016.
- O'Sullivan, P., Dankaerts, W., Burnett, A., Straker, L., Bargon, G., Moloney, N., Perry, M., and Tsang, S., 2006c. Lumbopelvic kinematics and trunk muscle activity during sitting on stable and unstable surfaces. *Journal of Orthopaedic & Sports Physical Therapy*, 36, 19–25.
- O'Sullivan, K., O'Dea, P., Dankaerts, W., O'Sullivan, P., Clifford, A., and O'Sullivan, L., 2010. Neutral lumbar spine sitting posture in pain-free subjects. *Manual Therapy*, 15, 557–561.
- O'Sullivan, P.B., Smith, A.J., Beales, D.J., and Straker, L.M., 2011a. Association of biopsychosocial factors with degree of slump in sitting posture and self-report of back pain in adolescents: a cross-sectional study. *Physical Therapy*, 91, 470–488.
- O'Sullivan, K., Galleotti, L., Dankaerts, W., O'Sullivan, P., and O'Sullivan, L., 2011b. The between-day and inter-rater reliability of a novel wireless system to analyse lumbar spine posture. *Ergonomics*, 54, 82–90.
- O'Sullivan, K., O'Sullivan, L., Campbell, A., O'Sullivan, P., and Dankaerts, W., 2012. Towards monitoring lumbo-pelvic posture in real-life situations: concurrent validity of a novel posture monitor and a traditional laboratory-based motion analysis system. *Manual Therapy*, 17, 77–83.
- Roffey, D.M., Wai, E.K., Bishop, P., Kwon, B.K., and Dagenais, S., 2010. Causal assessment of occupational sitting and low back pain: results of a systematic review. *The Spine Journal*, 10, 252–261.
- Scannell, J.P. and McGill, S.M., 2003. Lumbar posture—should it, and can it, be modified? A study of passive tissue stiffness and lumbar position during activities of daily living. *Physical Therapy*, 83, 907–917.
- Seah, S.H.H., Briggs, A.M., O'Sullivan, P.B., Smith, A.J., Burnett, A.F., and Straker, L.M., 2011. An exploration of familial associations in spinal posture defined using a clinical grouping method. *Manual Therapy*, 16, 501–509.
- Smith, A.J., O'Sullivan, P.B., Campbell, A.C., and Straker, L.M., 2010. The relationship between back muscle endurance and physical, lifestyle, and psychological factors in adolescents. *Journal of Orthopaedic & Sports Physical Therapy*, 40, 517–523.
- Smith, A., O'Sullivan, P., Beales, D., De Klerk, N., and Straker, L., 2011. Trajectories of childhood body mass index are associated with adolescent sagittal standing posture. *International Journal of Paediatric Obesity*, 6, e97–e106.
- Soderberg, G. and Knutson, L.M., 2000. A guide for the use and interpretation of kinesiological electromyographic data. *Physical Therapy*, 80, 485–498.
- van Deursen, L., Patijn, J., Durinck, J., Brouwer, R., van Erven-Sommers, J., and Vortman, B., 1999. Sitting and low back pain: the positive effect of rotatory dynamic stimuli during prolonged sitting. *European Spine Journal*, 8, 187–193.
- van Dieen, J., De Looze, M., and Hermans, V., 2001. Effects of dynamic office chairs on trunk kinematics, trunk extensor EMG and spinal shrinkage. *Ergonomics*, 44, 739–750.
- Vergara, M. and Page, A., 2000. System to measure the use of the backrest in sitting-posture office tasks. *Applied Ergonomics*, 31, 247–254.
- Vergara, M. and Page, A., 2002. Relationship between comfort and back posture and mobility in sitting-posture. *Applied Ergonomics*, 33, 1–8.
- Westgaard, R.H. and DeLuca, C.J., 1999. Motor unit substitution in long-duration contractions of the human trapezius muscle. *Journal of Neurophysiology*, 82, 501–504.
- Womersley, L. and May, S., 2006. Sitting posture of subjects with postural backache. *Journal of Manipulative and Physiological Therapeutics*, 29, 213–218.